Author index

Abe, J.P. *see* Chung, W.H.
Acharya, S.N. *see* Li, G.Q.
Agarwal, J. *see* Ramachandran, P.
Ahlawat, Y.S. *see* Ramachandran, P.
Ahoonmanesh, A. *see* Farzadfar, Sh.
Akino, S. *see* Gotoh, K.
Al Maqbali, Y. *see* Al-Bahry, S.
Al Raeesi, A. *see* Al-Bahry, S.
Al Sa'di, A. *see* Al-Bahry, S.
Al-Bahry, S., Elshafie, A.E., Deadman, M., Al Sa'di, A., Al Raeesi, A. & Al Maqbali, Y. First report of *Ganoderma colossum* on *Ficus altissima* and *Delonix regia* in Oman, 245
Ala-Poikela, M., Svensson, E., Rojas, A., Horko, T., Paulin, L., Valkonen, J.P.T. & Kvarnheden, A. Genetic diversity and mixed infections of begomoviruses infecting tomato, pepper and cucurbit crops in Nicaragua, 448
Alexander, H.M. *see* Thrall, P.H.
Almeida, R. *see* Arocha, Y.
Altınok, H.H. First report of fusarium wilt of eggplant caused by *Fusarium oxysporum* f. sp. *melongenae* in Turkey, 577
Alves-Santos, F.M. *see* Santamaría, O.
Amiri, A. & Bompeix, G. Diversity and population dynamics of *Penicillium* spp. on apples in pre- and postharvest environments: consequences for decay development, 74
Amrao, L. *see* Iram, S.
Andrivon, D. *see* Forbes, G.A.
see also Pasco, C.
see also Pilet, F.
Antesana, O. *see* Jones, P.
Anthony, F., Topart, P., Martinez, A., Silva, M. & Nicole, M. Hypersensitive-like reaction conferred by the *Mex-1* resistance gene against *Meloidogyne exigua* in coffee, 476
Anthony, K. *see* Bharathan, N.
Arabolaza, A.L. *see* Dunger, G.
Aramburu, J. & Galipienso, L. First report in Spain of a variant of *Tomato mosaic virus* (ToMV) overcoming the *Tm-2*^{2} resistance gene in tomato (*Lycopersicon esculentum*), 566
Aramburu, J. *see* Galipienso, L.
Arocha, Y., Horta, D., Piñol, B., Palenzuela, I., Picornell, S., Almeida, R. & Jones, P. First report of a phytoplasma associated with Bermuda-grass white leaf disease in Cuba, 233
Arocha, Y. *see* Jones, P.
Arocha, Y., López, M., Fernández, M., Piñol, B., Horta, D., Peralta, E.L., Almeida, R., Carvajal, O., Picornell, S., Wilson, M.R. & Jones, P. Transmission of a sugarcane yellow leaf phytoplasma by the delphacid planthopper *Saccharosydne saccharivora*, a new vector of sugarcane yellow leaf syndrome, 634
Arruda, G.M.T., Miller, R.N.G., Ferreira, M.A.S.V. & Café-Filho, A.C. Morphological and molecular characterization of the sudden-death syndrome pathogen of soybean in Brazil, 53
Auld, B.A. *see* Gilbert, R.L.
Awasthi, L.P. *see* Jain, R.K.
Aysan, Y. *see* Mirik, M.

Bag, S. *see* Jain, R.K.
Baloglu, S. *see* Mirik, M.
Bar-Joseph, M. *see* Gera, A.
Barba, M. *see* Tomassoli, L.
Barbara, D.J. *see* Collins, A.
Barnes, I., Nakabonge, G., Roux, J., Wingfield, B.D. & Wingfield, M.J. Comparison of populations of the wilt pathogen *Ceratocystis albifundus* in South Africa and Uganda, 189
Barreto, R.W. *see* Soares, D.J.
Barrett, L.G. *see* Thrall, P.H.
Basim, E. *see* Basim, H.
Basim, H., Basim, E., Yilmaz, S. & Ilkucan, M. First report of pith necrosis of tomato caused by *Pseudomonas mediterranea* in Turkey, 240
Bateman, G.L. *see* Freeman, J.
Bateman, G.L. The contribution of ground-level inoculum of *Fusarium culmorum* to ear blight of winter wheat, 428
Beales, P.A. *see* Lane, C.R.
Beasley, J.P. *see* Rowland, D.
Berg, T., Tesoriero, L. & Hailstones, D.L. PCR-based detection of *Xanthomonas campestris* pathovars in *Brassica* seed, 416
Bernal, A. *see* Díaz, J.
Bertaccini, A. *see* Rančić, D.
Bertrand, B. *see* Herve, G.
Bharathan, N., Saso, H., Gudipati, L., Bharathan, S., Whited, K. & Anthony, K. Double-stranded RNA: distribution and analysis among isolates of *Rhizoctonia solani* AG-2 to -13, 196
Bharathan, S. *see* Bharathan, N.
Blachinsky, D. *see* Kleitman, F.
Blechert, O. & Debener, T. Morphological characterization of the interaction between *Diplocarpon rosae* and various rose species, 82
Bloomer, P. *see* Coetzee, M.P.A.
Bock, C.H. *see* Wan, A.M.
Boland, G.J. *see* Kora, C.
Bompeix, G. *see* Amiri, A.
Bongiovanni, M. *see* Jacquet, M.
Boudon-Padieu, E. *see* Milkus, B.
Brading, P.A. *see* Chartrain, L.
Brasier, C.M. *see* Denman, S.
Brennan, J.M., Egan, D., Cooke, B.M. & Doohan, F.M. Effect of temperature on head blight of wheat caused by *Fusarium culmorum* and *F. graminearum*, 156
Briddon, R.W. *see* Iram, S.
Britz, H., Coutinho, T.A., Wingfield, B.D., Marasas, W.F.O. & Wingfield, M.J. Diversity and differentiation in two populations of *Gibberella circinata* in South Africa, 46
Brown, J.K.M. *see* Chartrain, L.
Budding, D.J. *see* Forbes, G.A.
Burdon, J.J. *see* Thrall, P.H.
Burr, T.J. *see* Kaewnum, S.
Buschmann, H., Gonsior, G. & Sauerborn, J. Pathogenicity of branched broomrape (*Orobanche ramosa*) populations on tobacco cultivars, 650

Café-Filho, A.C. *see* Arruda, G.M.T.
Caffier, V. & Laurens, F. Breakdown of *Pl2*, a major gene of resistance to apple powdery mildew, in a French experimental orchard, 116

© 2005 BSPP

Author index

Abe, J.P. *see* Chung, W.H.
Acharya, S.N. *see* Li, G.Q.
Agarwal, J. *see* Ramachandran, P.
Ahlawat, Y.S. *see* Ramachandran, P.
Ahoonmanesh, A. *see* Farzadfar, Sh.
Akino, S. *see* Gotoh, K.
Al Maqbali, Y. *see* Al-Bahry, S.
Al Raeesi, A. *see* Al-Bahry, S.
Al Sa'di, A. *see* Al-Bahry, S.
Al-Bahry, S., Elshafie, A.E., Deadman, M., Al Sa'di, A., Al Raeesi, A. & Al Maqbali, Y. First report of *Ganoderma colossum* on *Ficus altissima* and *Delonix regia* in Oman, 245
Ala-Poikela, M., Svensson, E., Rojas, A., Horko, T., Paulin, L., Valkonen, J.P.T. & Kvarnheden, A. Genetic diversity and mixed infections of begomoviruses infecting tomato, pepper and cucurbit crops in Nicaragua, 448
Alexander, H.M. *see* Thrall, P.H.
Almeida, R. *see* Arocha, Y.
Altınok, H.H. First report of fusarium wilt of eggplant caused by *Fusarium oxysporum* f. sp. *melongenae* in Turkey, 577
Alves-Santos, F.M. *see* Santamaría, O.
Amiri, A. & Bompeix, G. Diversity and population dynamics of *Penicillium* spp. on apples in pre- and postharvest environments: consequences for decay development, 74
Amrao, L. *see* Iram, S.
Andrivon, D. *see* Forbes, G.A.
see also Pasco, C.
see also Pilet, F.
Antesana, O. *see* Jones, P.
Anthony, F., Topart, P., Martinez, A., Silva, M. & Nicole, M. Hypersensitive-like reaction conferred by the *Mex-1* resistance gene against *Meloidogyne exigua* in coffee, 476
Anthony, K. *see* Bharathan, N.
Arabolaza, A.L. *see* Dunger, G.
Aramburu, J. & Galipienso, L. First report in Spain of a variant of *Tomato mosaic virus* (ToMV) overcoming the *Tm-2*^{2} resistance gene in tomato (*Lycopersicon esculentum*), 566
Aramburu, J. *see* Galipienso, L.
Arocha, Y., Horta, D., Piñol, B., Palenzuela, I., Picornell, S., Almeida, R. & Jones, P. First report of a phytoplasma associated with Bermuda-grass white leaf disease in Cuba, 233
Arocha, Y. *see* Jones, P.
Arocha, Y., López, M., Fernández, M., Piñol, B., Horta, D., Peralta, E.L., Almeida, R., Carvajal, O., Picornell, S., Wilson, M.R. & Jones, P. Transmission of a sugarcane yellow leaf phytoplasma by the delphacid planthopper *Saccharosydne saccharivora*, a new vector of sugarcane yellow leaf syndrome, 634
Arruda, G.M.T., Miller, R.N.G., Ferreira, M.A.S.V. & Café-Filho, A.C. Morphological and molecular characterization of the sudden-death syndrome pathogen of soybean in Brazil, 53
Auld, B.A. *see* Gilbert, R.L.
Awasthi, L.P. *see* Jain, R.K.
Aysan, Y. *see* Mirik, M.

Bag, S. *see* Jain, R.K.
Baloglu, S. *see* Mirik, M.
Bar-Joseph, M. *see* Gera, A.
Barba, M. *see* Tomassoli, L.
Barbara, D.J. *see* Collins, A.
Barnes, I., Nakabonge, G., Roux, J., Wingfield, B.D. & Wingfield, M.J. Comparison of populations of the wilt pathogen *Ceratocystis albifundus* in South Africa and Uganda, 189
Barreto, R.W. *see* Soares, D.J.
Barrett, L.G. *see* Thrall, P.H.
Basim, E. *see* Basim, H.
Basim, H., Basim, E., Yilmaz, S. & Ilkucan, M. First report of pith necrosis of tomato caused by *Pseudomonas mediterranea* in Turkey, 240
Bateman, G.L. *see* Freeman, J.
Bateman, G.L. The contribution of ground-level inoculum of *Fusarium culmorum* to ear blight of winter wheat, 428
Beales, P.A. *see* Lane, C.R.
Beasley, J.P. *see* Rowland, D.
Berg, T., Tesoriero, L. & Hailstones, D.L. PCR-based detection of *Xanthomonas campestris* pathovars in *Brassica* seed, 416
Bernal, A. *see* Díaz, J.
Bertaccini, A. *see* Rančić, D.
Bertrand, B. *see* Herve, G.
Bharathan, N., Saso, H., Gudipati, L., Bharathan, S., Whited, K. & Anthony, K. Double-stranded RNA: distribution and analysis among isolates of *Rhizoctonia solani* AG-2 to -13, 196
Bharathan, S. *see* Bharathan, N.
Blachinsky, D. *see* Kleitman, F.
Blechert, O. & Debener, T. Morphological characterization of the interaction between *Diplocarpon rosae* and various rose species, 82
Bloomer, P. *see* Coetzee, M.P.A.
Bock, C.H. *see* Wan, A.M.
Boland, G.J. *see* Kora, C.
Bompeix, G. *see* Amiri, A.
Bongiovanni, M. *see* Jacquet, M.
Boudon-Padieu, E. *see* Milkus, B.
Brading, P.A. *see* Chartrain, L.
Brasier, C.M. *see* Denman, S.
Brennan, J.M., Egan, D., Cooke, B.M. & Doohan, F.M. Effect of temperature on head blight of wheat caused by *Fusarium culmorum* and *F. graminearum*, 156
Briddon, R.W. *see* Iram, S.
Britz, H., Coutinho, T.A., Wingfield, B.D., Marasas, W.F.O. & Wingfield, M.J. Diversity and differentiation in two populations of *Gibberella circinata* in South Africa, 46
Brown, J.K.M. *see* Chartrain, L.
Budding, D.J. *see* Forbes, G.A.
Burdon, J.J. *see* Thrall, P.H.
Burr, T.J. *see* Kaewnum, S.
Buschmann, H., Gonsior, G. & Sauerborn, J. Pathogenicity of branched broomrape (*Orobanche ramosa*) populations on tobacco cultivars, 650

Café-Filho, A.C. *see* Arruda, G.M.T.
Caffier, V. & Laurens, F. Breakdown of *Pl2*, a major gene of resistance to apple powdery mildew, in a French experimental orchard, 116

© 2005 BSPP

Çakir, E. First report of potato wart disease in Turkey, 584
Capezio, S. *see* Forbes, G.A.
Carvajal, O. *see* Arocha, Y.
Castagnone-Sereno, P. *see* Jacquet, M.
Castillo, J. *see* Phillips-Mora, W.
Castillo, P. *see* Vovlas, N.
Cayford, B. *see* Saqib, M.
Cetinkaya-Yildiz, R. *see* Mirik, M.
Chacón, M.G. *see* Forbes, G.A.
Chang, Y.-C. *see* Lee, Y.-A.
see also Li, C.-Y.
Chartrain, L., Brading, P.A. & Brown, J.K.M. Presence of the *Stb6* gene for resistance to septoria tritici blotch (*Mycosphaerella graminicola*) in cultivars used in wheat-breeding programmes worldwide, 134
Chen, K.-P. *see* Lee, Y.-A.
Chen, S., Dai, X., Qiang, S. & Tang, Y. Effect of a nonhost-selective toxin from *Alternaria alternata* on chloroplast-electron transfer activity in *Eupatorium adenophorum*, 671
Chen, W. *see* Meng, X.
Chen, X.Y. *see* Zhu, X.Q.
Chhetri, D.B. *see* Coetzee, M.P.A.
Chow, S. *see* Fortune, M.P.
Chowda Reddy, R.V., Muniyappa, V., Colvin, J. & Seal, S. A new begomovirus isolated from *Gossypium barbadense* in southern India, 570
Chung, W.H., Abe, J.P., Yamaoka, Y., Haung, J.W. & Kakishima, M. First report of plumeria rust disease caused by *Coleosporium plumeriae* in Taiwan, 812
Ciampi, M.B. *see* Fenille, R.C.
Cilas, C. *see* Herve, G.
Ciuffo, M., Finetti-Sialer, M.M., Gallitelli, D. & Turina, M. First report in Italy of a resistance-breaking strain of *Tomato spotted wilt virus* infecting tomato cultivars carrying the *Sw5* resistance gene, 564
Ciuffo, M., Pettiti, D., Gallo, S., Masenga, V. & Turina, M. First report of *Blueberry scorch virus* in Europe, 565
Clair, D. *see* Milkus, B.
Clewes, E. *see* Collins, A.
Coetzee, M.P.A., Wingfield, B.D., Kirisits, T., Chhetri, D.B., Bloomer, P. & Wingfield, M.J. Identification of *Armillaria* isolates from Bhutan based on DNA sequence comparisons, 36
Collins, A., Mercado-Blanco, J., Jiménez-Díaz, R.M., Olivares, C., Clewes, E. & Barbara, D.J. Correlation of molecular markers and biological properties in *Verticillium dahliae* and the possible origins of some isolates, 549
Colon, L.T. *see* Forbes, G.A.
Colvin, J. *see* Chowda Reddy, R.V.
Constantinescu, O. *see* Lane, C.R.
Cooke, B.M. *see* Brennan, J.M.
Cooke, D.E.L. *see* Hussain, S.
Cooter, R.J. *see* Thresh, J.M.
Cosic, J. *see* Vrandecic, K.
Coutinho, T.A. *see* Britz, H.
Cuadrado, I.M. *see* Janssen, D.
Cunnington, J.H. *see* Pascoe, I.G.

Dai, X. *see* Chen, S.
Damsteegt, V.D. *see* Fatmi, M.
Das, D.R. *see* Ramachandran, P.
David, J. *see* Lane, C.R.
Davis, R.E. *see* Jomantiene, R.
De Dobbelaere, I. *see* De Jonghe, K.
De Jonghe, K., De Dobbelaere, I., Sarrazyn, R. & Höfte, M. Control of *Phytophthora cryptogea* in the hydroponic forcing of witloof chicory with the rhamnolipid-based biosurfactant formulation PRO1, 219
Deadman, M. *see* Al-Bahry, S.
Debener, T. *see* Blechert, O.
Delatte, H., Holota, H., Naze, F., Peterschmitt, M., Reynaud, B. & Lett, J.M. The presence of both recombinant and nonrecombinant strains of *Tomato yellow leaf curl virus* on tomato in Réunion Island, 262
deLittle, J.A. *see* Wilson, C.R.
Denman, S., Kirk, S.A., Brasier, C.M. & Webber, J.F. *In vitro* leaf inoculation studies as an indication of tree foliage susceptibility to *Phytophthora ramorum* in the UK, 512
Denman, S., Kirk, S.A., Brasier, C.M., Hughes, K.J.D., Griffin, R., Hobdon, E. & Webber, J.F. Foliar infection of sweet chestnut (*Castanea sativa*) by *Phytophthora ramorum* in the UK, 581
Denton, G. *see* Henricot, B.
Díaz, J., Silvar, C., Varela, M.M., Bernal, A. & Merino, F. *Fusarium* confers protection against several mycelial pathogens of pepper plants, 773
Diez, J.J. *see* Santamaría, O.
Dilbar, A. *see* Fortune, M.P.
Distel, S. *see* Forbes, G.A.
Dolby, C.A. *see* Stevens, M.
Doohan, F.M. *see* Brennan, J.M.
Dorner, J. *see* Rowland, D.
Doulis, A.G. *see* Vakalounakis, D.J.
Dror, O. *see* Kleitman, F.
Duan, C.J. *see* Li, Q.Q.
Duncan, J.M. *see* Forbes, G.A.
see also Hussain, S.
see also van de Graaf, P.
Dunemann, F. *see* Urbanietz, A.
Dunger, G., Arabolaza, A.L., Gottig, N., Orellano, E.G. & Ottado, J. Participation of *Xanthomonas axonopodis* pv. *citri hrp* cluster in citrus canker and nonhost plant responses, 781
Duvnjak, T. *see* Vrandecic, K.

Edison, S. *see* Jeeva, M.L.
Egan, D. *see* Brennan, J.M.
Elena, K. Fusarium wilt of *Phoenix canariensis*: first report in Greece, 244
Ellissèche, D. *see* Forbes, G.A.
see also Pilet, F.
Elmore, J. *see* Mumford, R.A.
Elshafie, A.E. *see* Al-Bahry, S.
Erickson, R.S. *see* Li, G.Q.
Evans, K.J., Jones, M.K. & Roush, R.T. Susceptibility of invasive taxa of European blackberry to rust disease caused by the uredinial stage of *Phragmidium violaceum* under field conditions in Australia, 275

Fang, R.X. *see* Sun, Q.H.
Farzadfar, Sh., Ohshima, K., Pourrahim, R., Golnaraghi, A.R., Jalali, S. & Ahoonmanesh, A. Occurrence of *Turnip mosaic virus* on ornamental crops in Iran, 261
Farzadfar, Sh., Pourrahim, R., Golnaraghi, A.R. & Ahoonmanesh, A. Occurrence of *Cauliflower mosaic virus* in different cruciferous plants in Iran, 810
Fatmi, M., Damsteegt, V.D. & Schaad, N.W. A combined agar-absorption and BIO-PCR assay for rapid, sensitive detection of *Xylella fastidiosa* in grape and citrus, 1

Feng, J.X. *see* Li, Q.Q.
Fenille, R.C., Ciampi, M.B., Souza, N.L., Nakatani, A.K. & Kuramae, E.E. Binucleate *Rhizoctonia* sp. AG G causing root rot in yacon (*Smallanthus sonchifolius*) in Brazil, 325
Fernández, M. *see* Arocha, Y.
Ferreira, M.A.S.V. *see* Arruda, G.M.T.
Finetti-Sialer, M.M. *see* Ciuffo, M.
Finlay, A.R. *see* Lane, C.R.
Fitt, B.D.L. *see* Wan, A.M.
Forbes, G.A., Chacón, M.G., Kirk, H.G., Huarte, M.A., Van Damme, M., Distel, S., Mackay, G.R., Stewart, H.E., Lowe, R., Duncan, J.M., Mayton, H.S., Fry, W.E., Andrivon, D., Ellissèche, D., Pellé, R., Platt, H.W., MacKenzie, G., Tarn, T.R., Colon, L.T., Budding, D.J., Lozoya-Saldaña, H., Hernandez-Vilchis, A. & Capezio, S. Stability of resistance to *Phytophthora infestans* in potato: an international evaluation, 364
Forbes, G.A. *see* Garry, G.
Fortune, M.P., Gosine, S., Chow, S., Dilbar, A., St. Hill, A., Gibbs, H. & Rambaran, N. First report of black sigatoka disease (causal agent *Mycosphaerella fijiensis*) from Trinidad, 246
Foulkes, M.J. *see* Pillinger, C.
Frampton, C.M. *see* McLear., K.L.
Franco, P. *see* Jones, P.
Frederickson, D.E. *see* Pažoutová, S.
Freeman, J., Ward, E., Gutteridge, R.J. & Bateman, G.L. Methods for studying population structure, including sensitivity to the fungicide silthiofam, of the cereal take-all fungus, *Gaeumannomyces graminis* var. *tritici*, 686
Fry, W.E. *see* Forbes, G.A.

Galipienso, L. *see* Aramburu, J.
Galipienso, L., Herranz, M.C., Pallás, V. & Aramburu, J. Detection of a tomato strain of *Parietaria mottle virus* (PMoV-T) by molecular hybridization and RT-PCR in field samples from north-eastern Spain, 29
Gallitelli, D. *see* Ciuffo, M.
Gallo, S. *see* Ciuffo, M.
García, B. *see* Velasco, L.
Garry, G., Forbes, G.A., Salas, A., Santa Cruz, M., Perez, W.G. & Nelson, R.J. Genetic diversity and host differentiation among isolates of *Phytophthora infestans* from cultivated potato and wild solanaceous hosts in Peru, 740
Ge, C. *see* Sun, Q.H.
Gefen, T. *see* Hadas, R.
Gera, A., Mawassi, M., Zeidan, M., Spiegel, S. & Bar-Joseph, M. An isolate of '*Candidatus* Phytoplasma australiense' group associated with Nivun Haamir dieback disease of papaya in Israel, 560
Gessler, C. *see* Gobbin, D.
Ghosh, D.K. *see* Ramachandran, P.
Gibb, K.S. *see* Streten, C.
Gibbs, H. *see* Fortune, M.P.
Gibson, L.J. *see* Wilson, C.R.
Gil, F. *see* Janssen, D.
Gilbert, R.L., Auld, B.A. & Hennecke, B.R. Leaf and stem spot of *Alternanthera philoxeroides* (alligatorweed) in Australia caused by *Nimbya* sp., 585
Gimeno, C. *see* Lannou, C.
Giorgi, S. & Scortichini, M. Molecular characterization of *Erwinia amylovora* strains from different host plants through RFLP analysis and sequencing of *hrp*N and *dsp*A/E genes, 789
Gobbin, D., Jermini, M., Loskill, B., Pertot, I., Raynal, M. & Gessler, C. Importance of secondary inoculum of *Plasmopara viticola* to epidemics of grapevine downy mildew, 522
Golnaraghi, A.R. *see* Farzadfar, S..
Gonsior, G. *see* Buschmann, H.
Gosine, S. *see* Fortune, M.P.
Gotoh, K., Akino, S., Maeda, A., Kondo, N., Naito, S., Kato, M. & Ogoshi, A. Characterization of some Asian isolates of *Phytophthora infestans*, 733
Gottig, N. *see* Dunger, G.
Goyeau, H. *see* Rimé, D.
Grau, C.R. *see* Meng, X.
Green, S. First report of *Septoria betulae* causing leaf spot of birch in the UK, 580
Griffin, R. *see* Denman, S.
Gryzenhout, M. *see* Rodas, C.A.
Gubler, W.D. *see* Rumbolz, J.
Gudipati, L. *see* Bharathan, N.
Guo, L.Y. *see* Zhu, X.Q.
Gutteridge, R.J. *see* Freeman, J.

Habili, N. *see* Milkus, B.
Hadas, R., Kritzman, G., Klietman, F., Gefen, T. & Manulis, S. Comparison of extraction procedures and determination of the detection threshold for *Clavibacter michiganensis* ssp. *michiganensis* in tomato seeds, 643
Hagiwara, K. *see* Kubo, C.
Haider, M.S. *see* Tahir, M.
Hailstones, D.L. *see* Berg, T.
Hallan, V. *see* Mehra, A.
see also Singh, A.K.
see also Verma, N.
Hallsworth, P.B. *see* Stevens, M.
Hansen, J.G., Koppel, M., Valskyte, A., Turka, I. & Kapsa, J. Evaluation of foliar resistance in potato to *Phytophthora infestans* based on an international field trial network, 169
Harju, V. *see* Mumford, R.A.
Harvey, J.L. *see* Wan, A.M.
Haung, J.W. *see* Chung, W.H.
He, C.Y. & Wolyn, D.J. Potential role for salicylic acid in induced resistance of asparagus roots to *Fusarium oxysporum* f.sp. *asparagi*, 227
He, C.Z. *see* Sun, Q.H.
Hegde, V. *see* Jeeva, M.L.
Hennecke, B.R. *see* Gilbert, R.L.
Henricot, B. & Denton, G. First record of the rust *Puccinia lagenophorae* on *Emilia* spp. in the UK, 242
Henricot, B. *see* Lane, C.R.
Hernandez-Vilchis, A. *see* Forbes, G.A.
Herranz, M.C. *see* Galipienso, L.
Herve, G., Bertrand, B., Villain, L., Licardie, D. & Cilas, C. Distribution analyses of *Meloidogyne* spp. and *Pratylenchus coffeae sensu lato* in coffee plots in Costa Rica and Guatemala, 471
Hiruki, C. *see* Wang, K.
Hita, I. *see* Velasco, L.
Hobdon, E. *see* Denman, S.
Höfte, M. *see* De Jonghe, K.
Holota, H. *see* Delatte, H.
Horko, T. *see* Ala-Poikela, M.
Horta, D. *see* Arocha, Y.
Hsiang, T. *see* Huang, J.
Hu, J. *see* Sun, Q.H.
Huang, G.X. *see* Sun, Q.H.

Huang, H.C. *see* Li, G.Q.
Huang, J., Zheng, L. & Hsiang, T. First report of leaf spot caused by *Curvularia verruculosa* on *Cynodon* sp. in Hubei, China, 253
Huarte, M.A. *see* Forbes, G.A.
Hubert, P. *see* Lannou, C.
Hughes, K.J.D. *see* Denman, S.
Hunt, J.S. *see* McLean, K.L.
Huo, N.X. *see* Li, A.L.
Hussain, D. *see* Rahman, M.
Hussain, S., Lees, A.K., Duncan, J.M. & Cooke, D.E.L. Development of a species-specific and sensitive detection assay for *Phytophthora infestans* and its application for monitoring of inoculum in tubers and soil, 373

Ichiki, T.U. *see* Kubo, C.
Idir, S. *see* Milkus, B.
Ilkucan, M. *see* Basim, H.
Iram, S., Amrao, L., Mansoor, M.S., Malik, A.H., Briddon, R.W. & Zafar, Y. First report of a begomovirus associated with leaf curl disease of *Duranta erecta* in Pakistan, 260

Jönsson, U., Jung, T., Sonesson, K. & Rosengren, U. Relationships between health of *Quercus robur*, occurrence of *Phytophthora* species and site conditions in southern Sweden, 502
Jacquet, M., Bongiovanni, M., Martinez, M., Verschave, P., Wajnberg, E. & Castagnone-Sereno, P. Variation in resistance to the root-knot nematode *Meloidogyne incognita* in tomato genotypes bearing the *Mi* gene, 93
Jain, R.K., Bag, S. & Awasthi, L.P. First report of natural infection of *Capsicum annuum* by *Tobacco streak virus* in India, 257
Jalali, S. *see* Farzadfar, S..
Janssen, D., Saez, E., Segundo, E., Martín, G., Gil, F. & Cuadrado, I.M. *Capsicum annuum* – a new host of *Parietaria mottle virus* in Spain, 567
Jarvis, B. *see* Mumford, R.A.
Jeeva, M.L., Hegde, V., Makeshkumar, T., Nair, R.R. & Edison, S. *Dioscorea alata*, a new host of *Sclerotium rolfsii* in India, 574
Jenkyn, J.F. *see* Wan, A.M.
Jermini, M. *see* Gobbin, D.
Jia, J.Z. *see* Li, A.L.
Jiménez-Díaz, R.M. *see* Collins, A.
Jomantiene, R. & Davis, R.E. Apple sessile leaf: a new disease associated with a '*Candidatus* Phytoplasma asteris' subgroup 16SrI-B phytoplasma in Lithuania, 237
Jones, M.G.K. *see* Saqib, M.
Jones, M.K. *see* Evans, K.J.
Jones, P. *see* Arocha, Y.
Jones, P., Arocha, Y., Antesana, O., Montellano, E. & Franco, P. 'Brotes grandes' (big bud) of potato: a new disease associated with a 16SrI-B subgroup phytoplasma in Bolivia, 234
Jones, P., Arocha, Y., Antesana, O., Montilliano, E. & Franco, P. First report of an isolate of '*Candidatus* Phytoplasma australiense' associated with a yellow leaf roll disease of peach (*Prunus persicae*) in Bolivia, 558
Jones, P., Arocha, Y., Antesana, O., Montellano, E. & Franco, P. 'Hoja de perejil' (parsley leaf) of tomato and morrenia little leaf, two new diseases associated with a phytoplasma in Bolivia, 235
Jones, P., Arocha, Y. & Plata, G. First report of a '*Candidatus* Phytoplasma asteris' isolate associated with a witches' broom disease of alfalfa in Bolivia, 559
Jones, R.A.C. *see* Saqib, M.
Jouan, B. *see* Pasco, C.
Jung, H.S. *see* Lee, H.B.
Jung, T. *see* Jönsson, U.
Jurkovic, D. *see* Vrandecic, K.

Kaewnum, S., Prathuangwong, S. & Burr, T.J. Aggressiveness of *Xanthomonas axonopodis* pv. *glycines* isolates to soybean and hypersensitivity responses by other plants, 409
Kajihara, H. *see* Kubo, C.
Kakishima, M. *see* Chung, W.H.
Kapsa, J. *see* Hansen, J.G.
Kato, M. *see* Gotoh, K.
Khan, A.A., Naqvi, Q.A., Khan, M.S., Singh, R. & Raj, S.K. First report of a begomovirus infecting *Calendula* in India, 569
Khan, M.S. *see* Khan, A.A.
see also Raj, S.K.
Kirisits, T. *see* Coetzee, M.P.A.
Kirk, H.G. *see* Forbes, G.A.
Kirk, S.A. *see* Denman, S.
Kitkaru, A.S. *see* Mantri, N.L.
Klietman, F. *see* Hadas, R.
Kleitman, F., Shtienberg, D., Blachinsky, D., Oppenheim, D., Zilberstaine, M., Dror, O. & Manulis, S. *Erwinia amylovora* populations resistant to oxolinic acid in Israel: prevalence, persistence and fitness, 108
Klironomou, E. *see* Vakalounakis, D.J.
Kondo, N. *see* Gotoh, K.
Kong, X.Y. *see* Li, A.L.
Koppel, M. *see* Hansen, J.G.
Kora, C., McDonald, M.R. & Boland, G.J. Occurrence of fungal pathogens of carrots on wooden boxes used for storage, 665
Kovač, M. *see* Krečič-Stres, H.
Krauss, U. *see* Phillips-Mora, W.
Krečič-Stres, H., Vučak, C., Ravnikar, M. & Kovač, M. Systemic *Potato virus* Y^{NTN} infection and levels of salicylic and gentisic acids in different potato genotypes, 441
Krishna Kishore, G., Pande, S. & Podile, A.R. Management of late leaf spot of groundnut (*Arachis hypogaea*) with chlorothalonil-tolerant isolates of *Pseudomonas aeruginosa*, 401
Kritzman, G. *see* Hadas, R.
Kubo, C., Nakazono-Nagaoka, E., Hagiwara, K., Kajihara, H., Takeuchi, S., Matsuo, K., Ichiki, T.U. & Omura, T. New severe strains of *Melon necrotic spot virus*: symptomatology and sequencing, 615
Kucera, L. *see* Leisova, L.
Kulshrestha, S. *see* Verma, N.
Kuramae, E.E. *see* Fenille, R.C.
Kurt, S. *see* Soylu, S.
Kurt, S., Soylu, E.M., Soylu, S. & Tok, F.M. First report of crater rot caused by *Rhizoctonia carotae* on carrots (*Daucus carota*) in Turkey, 251
Kusek, M. *see* Mirik, M.
Kvarnheden, A. *see* Ala-Poikela, M.

Lal, B. *see* Mehra, A.
Landa, B.B. *see* Vovlas, N.
Lane, C.R., Beales, P.A., O'neill, T.M., McPherson, G.M., Finlay, A.R., David, J., Constantinescu, O. & Henricot, B. First report of *Impatiens* downy mildew (*Plasmopara obducens*) in the UK, 243
Lannou, C., Hubert, P. & Gimeno, C. Competition and interactions among stripe rust pathotypes in wheat-cultivar mixtures, 699

Lannou, C. *see* Rimé, D.
Laurens, F. *see* Caffier, V.
Lee, H.B., Park, J.Y. & Jung, H.S. First report of leaf anthracnose caused by *Colletotrichum boninense* on spindle trees, 254
Lee, Y.-A., Chen, K.-P. & Chang, Y.-C. First report of bacterial leaf blight of white-flowered calla lily caused by *Xanthomonas campestris* pv. *zantedeschiae* in Taiwan, 239
Lees, A.K. *see* Hussain, S.
see also van de Graaf, P.
Leisova, L., Kucera, L., Minarikova, V. & Ovesna, J. AFLP-based PCR markers that differentiate spot and net forms of *Pyrenophora teres*, 66
Lemaire, O. *see* Stevens, M.
Lett, J.M. *see* Delatte, H.
Li, A.L., Wang, M.L., Zhou, R.H., Kong, X.Y., Huo, N.X., Wang, W.S. & Jia, J.Z. Comparative analysis of early H_2O_2 accumulation in compatible and incompatible wheat–powdery mildew interactions, 308
Li, B.-D. *see* Li, B.-H.
Li, B.-H., Xu, X.-M., Li, J.-T. & Li, B.-D. Effects of temperature and continuous and interrupted wetness on the infection of pear leaves by conidia of *Venturia nashicola*, 357
Li, C.-Y. & Chang, Y.-C. First identification of *Tobacco mild green mosaic virus* on *Capsicum annuum* in Taiwan, 258
Li, G.Q., Huang, H.C., Acharya, S.N. & Erickson, R.S. Effectiveness of *Coniothyrium minitans* and *Trichoderma atroviride* in suppression of sclerotinia blossom blight of alfalfa, 204
Li, J.-T. *see* Li, B.-H.
Li, Q.Q., Feng, J.X., Tang, J.L., Lin, W., Duan, C.J., Ye, Y.F. & Luo, K. *Siraita grosvenorii* (Luo Han Guo; Cucurbitaceae) is a new host of *Ralstonia solanacearum* in China, 811
Liang, Y.C., Sun, W.C., Si, J. & Römheld, V. Effects of foliar- and root-applied silicon on the enhancement of induced resistance to powdery mildew in *Cucumis sativus*, 678
Licardie, D. *see* Herve, G.
Lin, W. *see* Li, Q.Q.
López, M. *see* Arocha, Y.
Loskill, B. *see* Gobbin, D.
Lowe, R. *see* Forbes, G.A.
Lozoya-Saldaña, H. *see* Forbes, G.A.
Luo, K. *see* Li, Q.Q.
Luo, Y. *see* Zhu, X.Q.

Małolepsza, U. Spatial and temporal variation of reactive oxygen species and antioxidant enzymes in *o*-hydroxyethylorutin-treated tomato leaves inoculated with *Botrytis cinerea*, 317
Mackay, G.R. *see* Forbes, G.A.
MacKenzie, G. *see* Forbes, G.A.
Mackey, I.C. *see* Tzanetakis, I.E.
Maeda, A. *see* Gotoh, K.
Maiti, S. *see* Mandal, K.
Makeshkumar, T. *see* Jeeva, M.L.
Malathi, V.G. *see* Rouhibakhsh, A.
Malik, A.H. *see* Iram, S.
Malik, T.A. *see* Rahman, M.
Mandal, K. & Maiti, S. Bacterial soft rot of aloe caused by *Pectobacterium chrysanthemi*: a new report from India, 573
Mansoor, M.S. *see* Iram, S.
Mantri, N.L., Kitkaru, A.S., Misal, M.B. & Ravi, K.S. First report of *Papaya ringspot virus* – W in bottle gourd (*Lagenaria siceraria*) from India, 806
Manulis, S. *see* Hadas, R.
see also Kleitman, F.
Marasas, W.F.O. *see* Britz, H.
Martín, G. *see* Janssen, D.
Martin, R.R. *see* Tzanetakis, I.E.
Martinez, A. *see* Anthony, F.
Martinez, M. *see* Jacquet, M.
Masenga, V. *see* Ciuffo, M.
Matsuo, K. *see* Kubo, C.
Mawassi, M. *see* Gera, A.
Mayton, H.S. *see* Forbes, G.A.
McDonald, M.R. *see* Kora, C.
McLean, K.L., Swaminathan, J., Frampton, C.M., Hunt, J.S., Ridgway, H.J. & Stewart, A. Effect of formulation on the rhizosphere competence and biocontrol ability of *Trichoderma atroviride* C52, 212
McPherson, G.M. *see* Lane, C.R.
Mehra, A., Hallan, V., Lal, B., Negi, H.R. & Zaidi, A.A. *Sorbaria tomentosa* – a new natural host for *Potato virus Y* (PVY), 572
Meng, X., Grau, C.R. & Chen, W. Cultivar preference exhibited by two sympatric and genetically distinct populations of the soybean fungal pathogen *Phialophora gregata* f.sp. *sojae*, 180
Mercado-Blanco, J. *see* Collins, A.
Merino, F. *see* Díaz, J.
Mifsud, D. *see* Vovlas, N.
Milkus, B., Clair, D., Idir, S., Habili, N. & Boudon-Padieu, E. First detection of stolbur phytoplasma in grapevines (*Vitis vinifera* cv. Chardonnay) affected with grapevine yellows in the Ukraine, 236
Miller, R.N.G. *see* Arruda, G.M.T.
Minarikova, V. *see* Leisova, L.
Mirik, M., Baloglu, S., Aysan, Y., Cetinkaya-Yildiz, R., Kusek, M. & Sahin, F. First outbreak and occurrence of citrus blast disease, caused by *Pseudomonas syringae* pv. *syringae*, on orange and mandarin trees in Turkey, 238
Misal, M.B. *see* Mantri, N.L.
Montellano, E. *see* Jones, P.
Mumford, R.A., Jarvis, B., Harju, V., Elmore, J. & Skelton, A. The first identification of two viruses infecting trailing verbena in the UK, 568
Muniyappa, V. *see* Chowda Reddy, R.V.
Myburg, H. *see* Rodas, C.A.

Nair, R.R. *see* Jeeva, M.L.
Naito, S. *see* Gotoh, K.
Nakabonge, G. *see* Barnes, I.
Nakatani, A.K. *see* Fenille, R.C.
Nakazono-Nagaoka, E. *see* Kubo, C.
Naqvi, Q.A. *see* Khan, A.A.
Naze, F. *see* Delatte, H.
Negi, H.R. *see* Mehra, A.
Nelson, R.J. *see* Garry, G.
Nicole, M. *see* Anthony, F.
Nikolaou, K. *see* Tsopelas, P.

Ogoshi, A. *see* Gotoh, K.
Ohshima, K. *see* Farzadfar, S..
Olivares, C. *see* Collins, A.
Omura, T. *see* Kubo, C.
O'Neill, T.M. *see* Lane, C.R.
Oppenheim, D. *see* Kleitman, F.
Orellano, E.G. *see* Dunger, G.
Oros, G. *see* Vajna, L.

Ottado, J. *see* Dunger, G.
Ovesna, J. *see* Leisova, L.

Padilla, V. *see* Velasco, L.
Palenzuela, I. *see* Arocha, Y.
Pallás, V. *see* Galipienso, L.
Paltrinieri, S. *see* Rančić, D.
Pande, S. *see* Krishna Kishore, G.
Park, J.Y. *see* Lee, H.B.
Pasco, C., Jouan, B. & Andrivon, D. Resistance of potato genotypes to common and netted scab-causing species of *Streptomyces*, 383
Pascoe, I.G., Priest, M.J., Shivas, R.G. & Cunnington, J.H. Ustilospores of *Tilletia ehrhartae*, a smut of *Ehrharta calycina*, are common contaminants of Australian wheat grain, and a potential source of confusion with *Tilletia indica*, the cause of Karnal bunt of wheat, 161
Patron, N.J. *see* Stevens, M.
Paulin, L. *see* Ala-Poikela, M.
Paveley, N. *see* Pillinger, C.
Pažoutová, S. & Frederickson, D.E. Genetic diversity of *Claviceps africana* on sorghum and *Hyparrhenia*, 749
Pellé, R. *see* Forbes, G.A.
see also Pilet, F.
Peralta, E.L. *see* Arocha, Y.
Perez, W.G. *see* Garry, G.
Pertot, I. *see* Gobbin, D.
Petanović, R. *see* Rančić, D.
Peterschmitt, M. *see* Delatte, H.
Pettiti, D. *see* Ciuffo, M.
Phillips-Mora, W., Castillo, J., Krauss, U., Rodríguez, E. & Wilkinson, M.J. Evaluation of cacao (*Theobroma cacao*) clones against seven Colombian isolates of *Moniliophthora roreri* from four pathogen genetic groups, 483
Piątek, M. First report of powdery mildew of *Ligustrum* caused by *Erysiphe syringae-japonicae*, 578
Picornell, S. *see* Arocha, Y.
Pilet, F., Pellé, R., Ellissèche, D. & Andrivon, D. Efficacy of the *R2* resistance gene as a component for the durable management of potato late blight in France, 723
Pillinger, C., Paveley, N., Foulkes, M.J. & Spink, J. Explaining variation in the effects of take-all (*Gaeumannomyces graminis* var. *tritici*) on nitrogen and water uptake by winter wheat, 491
Piñol, B. *see* Arocha, Y.
Plata, G. *see* Jones, P.
Platt, H.W. *see* Forbes, G.A.
Podile, A.R. *see* Krishna Kishore, G.
Pourrahim, R. *see* Farzadfar, Sh.
Prathuangwong, S. *see* Kaewnum, S.
Priest, M.J. *see* Pascoe, I.G.
Putnam, M.L. *Allantophomopsis lycopodina* – a new aerial pathogen of lingonberry (*Vaccinium vitis-idaea*), 248

Qiang, S. *see* Chen, S.

Rahman, M., Hussain, D., Malik, T.A. & Zafar, Y. Genetics of resistance to cotton leaf curl disease in *Gossypium hirsutum*, 764
Raikhy, G. *see* Verma, N.
Raj, S.K., Khan, M.S. & Singh, R. Natural occurrence of a begomovirus on pigeonpea in India, 809
Raj, S.K. *see* Khan, A.A.
Ram, R. *see* Singh, A.K.
see also Verma, N.
Ramachandran, P., Agarwal, J., Roy, A., Ghosh, D.K., Das, D.R. & Ahlawat, Y.S. First record of a *Hop stunt viroid* variant on Nagpur mandarin and Mosambi sweet orange trees on rough lemon and Rangpur lime rootstocks, 571
Rambaran, N. *see* Fortune, M.P.
Rančić, D., Paltrinieri, S., Toševski, I., Petanović, R., Stevanović, B. & Bertaccini, A. First report of multiple inflorescence disease of *Cirsium arvense* and its association with a 16SrIII-B subgroup phytoplasma in Serbia, 561
Ravi, K.S. *see* Mantri, N.L.
Ravnikar, M. *see* Krečič-Stres, H.
Raynal, M. *see* Gobbin, D.
Reynaud, B. *see* Delatte, H.
Riccioni, L. *see* Vrandecic, K.
Ridgway, H.J. *see* McLean, K.L.
Rimé, D., Robert, C., Goyeau, H. & Lannou, C. Effect of host genotype on leaf rust (*Puccinia triticina*) lesion development and urediniospore production in wheat seedlings, 287
Robert, C. *see* Rimé, D.
Robinson, J. *see* Xu, X.M.
Rodas, C.A., Gryzenhout, M., Myburg, H., Wingfield, B.D. & Wingfield, M.J. Discovery of the *Eucalyptus* canker pathogen *Chrysoporthe cubensis* on native *Miconia* (Melastomataceae) in Colombia, 460
Rodríguez, E. *see* Phillips-Mora, W.
Rojas, A. *see* Ala-Poikela, M.
Römheld, V. *see* Liang, Y.C.
Ronco, L. *see* Sisterna, M.
Rosengren, U. *see* Jönsson, U.
Rouhibakhsh, A. & Malathi, V.G. Severe leaf curl disease of cowpea – a new disease of cowpea in northern India caused by *Mungbean yellow mosaic India virus* and a satellite DNA β, 259
Roush, R.T. *see* Evans, K.J.
Roux, J. *see* Barnes, I.
Rowland, D., Dorner, J., Sorensen, R., Beasley, J.P. & Todd, J. *Tomato spotted wilt virus* in peanut tissue types and physiological effects related to disease incidence and severity, 431
Roy, A. *see* Ramachandran, P.
Rumbolz, J. & Gubler, W.D. Susceptibility of grapevine buds to infection by powdery mildew *Erysiphe necator*, 535

Saez, E. *see* Janssen, D.
Safrankova, I. First report of *Volutella pachysandricola* on *Pachysandra terminalis* in the Czech Republic, 808
Sahin, F. *see* Mirik, M.
Salas, A. *see* Garry, G.
Santa Cruz, M. *see* Garry, G.
Santamaría, O., Alves-Santos, F.M. & Diez, J.J. Genetic characterization of *Gremmeniella abietina* var. *abietina* isolates from Spain, 331
Saqib, M., Jones, R.A.C., Cayford, B. & Jones, M.G.K. First report of *Bean common mosaic virus* in Western Australia, 563
Sarrazyn, R. *see* De Jonghe, K.
Saso, H. *see* Bharathan, N.
Sauerborn, J. *see* Buschmann, H.
Schaad, N.W. *see* Fatmi, M.
Schlenzig, A. First report of *Phytophthora inflata* on nursery plants of *Rhododendron* spp., *Gaultheria shalon* and *Vaccinium vitis-idaea* in Scotland, 582
Schupp, P.J. *see* Wilson, C.R.
Scortichini, M. *see* Giorgi, S.

Seal, S. *see* Chowda Reddy, R.V.
Segundo, E. *see* Janssen, D.
Sert, H.B. & Sümbül, H. First report of leaf spot caused by *Phoma sorghina* on *Trifolium campestre* in Turkey, 249
Sert, H.B. & Sümbül, H. First report of *Melampsoridium hiratsukanum* infecting alder (*Alnus orientalis* var. *orientalis*) in Turkey, 241
Shivas, R.G. *see* Pascoe, I.G.
Shtienberg, D. *see* Kleitman, F.
Si, J. *see* Liang, Y.C.
Silva, M. *see* Anthony, F.
Silvar, C. *see* Díaz, J.
Singh, A.K., Hallan, V., Ram, R. & Zaidi, A.A. Variability in the coat protein of *Lily symptomless virus* isolates infecting various lily species, 621
Singh, A.K. *see* Verma, N.
Singh, A.K., Hallan, V., Verma, N. & Zaidi, A.A. Natural occurrence of *Lily symptomless virus* on spider lily, 255
Singh, L. *see* Verma, N.
Singh, R. *see* Khan, A.A.
see also Raj, S.K.
Sisterna, M. & Ronco, L. Occurrence of grapevine leaf spot caused by *Pseudocercospora vitis* in Argentina, 247
Skelton, A. *see* Mumford, R.A.
Smith, H.G. *see* Stevens, M.
Soares, D.J. & Barreto, R.W. First report of *Oidium piperis* on *Piper aduncum* in Brazil, 579
Sonesson, K. *see* Jönsson, U.
Sorensen, R. *see* Rowland, D.
Souza, N.L. *see* Fenille, R.C.
Soylu, E.M. *see* Kurt, S.
see also Soylu, S.
Soylu, S., Kurt, S., Soylu, E.M. & Tok, F.M. First report of alternaria leaf blight caused by *Alternaria dauci* on carrot in Turkey, 252
Soylu, S. *see* Kurt, S.
Spiegel, S. *see* Gera, A.
Spink, J. *see* Pillinger, C.
St. Hill, A. *see* Fortune, M.P.
Stead, D.E. *see* Weller, S.A.
Stevanović, B. *see* Rančić, D.
Stevens, M., Patron, N.J., Dolby, C.A., Weekes, R., Hallsworth, P.B., Lemaire, O. & Smith, H.G. Distribution and properties of geographically distinct isolates of sugar beet yellowing viruses, 100
Stewart, A. *see* McLean, K.L.
Stewart, H.E. *see* Forbes, G.A.
Streten, C. & Gibb, K.S. Genetic variation in *Candidatus* Phytoplasma australiense, 8
Sümbül, H. *see* Sert, H.B.
Sun, Q.H., Hu, J., Huang, G.X., Ge, C., Fang, R.X. & He, C.Z. Type-II secretion pathway structural gene *xpsE*, xylanase- and cellulase secretion and virulence in *Xanthomonas oryzae* pv. *oryzae*, 15
Sun, W.C. *see* Liang, Y.C.
Svensson, E. *see* Ala-Poikela, M.
Swaminathan, J. *see* McLean, K.L.

Tahir, M. & Haider, M.S. First report of *Tomato leaf curl New Delhi virus* infecting bitter gourd in Pakistan, 807
Takeuchi, S. *see* Kubo, C.
Tang, J.L. *see* Li, Q.Q.
Tang, Y. *see* Chen, S.
Tarn, T.R. *see* Forbes, G.A.
Tesoriero, L. *see* Berg, T.
Thrall, P.H., Barrett, L.G., Burdon, J.J. & Alexander, H.M. Variation in pathogen aggressiveness within a metapopulation of the *Cakile maritima–Alternaria brassicicola* host–pathogen association, 265
Thresh, J.M. & Cooter, R.J. Strategies for controlling cassava mosaic virus disease in Africa, 587
Toševski, I. *see* Rančić, D.
Todd, J. *see* Rowland, D.
Tok, F.M. *see* Kurt, S.
see also Soylu, S.
Tomassoli, L., Zaccaria, A. & Barba, M. *Capparis spinosa* – a new host of *Cucumber mosaic virus* in Italy, 263
Topart, P. *see* Anthony, F.
Tsopelas, P. & Nikolaou, K. First report of *Heterobasidion annosum* in Cyprus, 583
Turina, M. *see* Ciuffo, M.
Turka, I. *see* Hansen, J.G.
Tzanetakis, I.E., Mackey, I.C. & Martin, R.R. *Tulip virus X* (TVX) associated with lemon balm (*Melissa officinalis*) variegation: first report of TVX in the USA, 562

Urbanietz, A. & Dunemann, F. Isolation, identification and molecular characterization of physiological races of apple powdery mildew (*Podosphaera leucotricha*), 125

Vajna, L. & Oros, G. First report of *Rhizoctonia zeae* on *Lolium perenne* and *Festuca* sp. in Hungary, 250
Vakalounakis, D.J., Doulis, A.G. & Klironomou, E. Characterization of *Fusarium oxysporum* f. sp. *radicis-cucumerinum* attacking melon under natural conditions in Greece, 339
Valkonen, J.P.T. *see* Ala-Poikela, M.
Valskyte, A. *see* Hansen, J.G.
Van Damme, M. *see* Forbes, G.A.
van de Graaf, P., Lees, A.K., Wale, S.J. & Duncan, J.M. Effect of soil inoculum level and environmental factors on potato powdery scab caused by *Spongospora subterranea*, 22
Varela, M.M. *see* Díaz, J.
Velasco, L., García, B., Hita, I. & Padilla, V. Detection of *Grapevine virus B* associated with rugose wood (corky bark) symptoms in grapevine cv. Napoleon in Murcia (Spain), 264
Verma, N., Singh, L., Singh, A.K., Kulshrestha, S., Raikhy, G., Hallan, V., Ram, R. & Zaidi, A.A. Ornithogalum: a new host of *Cucumber mosaic virus* (CMV) from India, 256
Verma, N. *see* Singh, A.K.
Verschave, P. *see* Jacquet, M.
Villain, L. *see* Herve, G.
Vovlas, N., Mifsud, D., Landa, B.B. & Castillo, P. Pathogenicity of the root-knot nematode *Meloidogyne javanica* on potato, 657
Vrandecic, K., Cosic, J., Riccioni, L., Duvnjak, T. & Jurkovic, D. Isolation of *Diaporthe phaseolorum* var. *caulivora* from *Abutilon theophrasti* in Croatia, 576
Vučak, C. *see* Krečič-Stres, H.

Wajnberg, E. *see* Jacquet, M.
Wale, S.J. *see* van de Graaf, P.
Wale, S.J. The science of appliance, 715
Wan, A.M., Bock, C.H., Fitt, B.D.L., Harvey, J.L. & Jenkyn, J.F. Development of *Oculimacula yallundae* and *O. acuformis* (eyespot) on leaf sheaths of winter wheat in the UK in relation to thermal time, 144
Wang, K. & Hiruki, C. Distinctions between phytoplasmas at the subgroup level detected by heteroduplex mobility assay, 625

Wang, M.L. *see* Li, A.L.
Wang, W.S. *see* Li, A.L.
Ward, E. *see* Freeman, J.
Webber, J.F. *see* Denman, S.
Weekes, R. *see* Stevens, M.
Weller, S.A., Stead, D.E. & Young, J.P.W. Induction of root-mat symptoms on cucumber plants by *Rhizobium*, but not by *Ochrobactrum* or *Sinorhizobium*, harbouring a cucumopine Ri plasmid, 799
Whited, K. *see* Bharathan, N.
Wilkinson, M.J. *see* Phillips-Mora, W.
Wilson, C.R., deLittle, J.A., Wong, J.A.L., Schupp, P.J. & Gibson, L.J. Adjustment of soil-surface pH and comparison with conventional fungicide treatments for control of lettuce drop (*Sclerotinia minor*), 393
Wilson, M.R. *see* Arocha, Y.
Wingfield, B.D. *see* Barnes, I.
see also Britz, H.
see also Coetzee, M.P.A.
see also Rodas, C.A.
Wingfield, M.J. *see* Barnes, I.
see also Britz, H.
see also Coetzee, M.P.A.
see also Rodas, C.A.
Wolyn, D.J. *see* He, C.Y.
Wong, J.A.L. *see* Wilson, C.R.

Xu, X.M. & Robinson, J. Modelling the effects of wetness duration and fruit maturity on infection of apple fruits of Cox's Orange Pippin and two clones of Gala by *Venturia inaequalis*, 347
Xu, X.-M. *see* Li, B.-H.

Yamaoka, Y. *see* Chung, W.H.
Ye, Y.F. *see* Li, Q.Q.
Yilmaz, S. *see* Basim, H.
Young, J.P.W. *see* Weller, S.A.

Zaccaria, A. *see* Tomassoli, L.
Zafar, Y. *see* Iram, S.
see also Rahman, M.
Zaidi, A.A. *see* Mehra, A.
see also Singh, A.K.
see also Verma, N.
Zeidan, M. *see* Gera, A.
Zheng, L. *see* Huang, J.
Zhou, R.H. *see* Li, A.L.
Zhu, X.Q., Chen, X.Y., Luo, Y. & Guo, L.Y. First report of *Monilinia fructicola* on peach and nectarine in China, 575
Zilberstaine, M. *see* Kleitman, F.

Keyword index

16/23S spacer region, 625
16S rRNA gene, 8
α-Proteobacteria, 799

Acacia mearnsii, 189
accumulated temperature, 144
AFLP, 339
AFLP marker analysis, 125
age-related susceptibility, 347
aggressiveness, 287, 723
airborne *Penicillium*, 74
alfalfa blossom blight, 204
allozyme analysis, 733
Amelanchier, 789
amphihaploids, 549
anastomosis group, 196
anastomosis grouping, 325
antibiotic resistance, 108
apple, 108
apple cultivars, 74
apple scab, 347
Arachis hypogaea, 401, 431
area under disease progress curve, 364
Armillaria mellea, 36
Armillaria root rot, 36
Asia, 36
attachment, 799
Auchenorrhyncha, 634

bacterial canker, 643
bacterial wilt, 643
barley net blotch, 66
begomovirus, 587
begomoviruses, 764
Bemisia tabaci, 587
Bhutan, 36
biodiversity, 625
biological control, 204, 212, 219
biological species, 36
BIO-PCR, 1
biotype, 331
black rot, 416
black spot, 82
blackberry rust, 275
blue mould, 74
Botrytis cinerea, 665
Brassica spp., 416
broad-leaved species, 512
brown stem rot, 180

$Ca(OH)_2$, 393
cacao, 483
$CaCO_3$, 393
calcium hydroxide, 393
Capsicum spp., 448
cassava, 587
Ceratocystis fimbriata, 189
Cerebella andropogonis, 749
chitinase, 773
Chrysoporthe cubensis, 460
Chrysoporthella hodgesiana, 460
Cichorium intybus var. *foliosum*, 219
citrus variegated chlorosis, 1
Claviceps spp., 749
clonality, 749
closterovirus, 100
cocoa, 483
Coffea arabica, 471, 476
Colombia, 460
compatible interaction, 308
competition, 471
competitive bioassay, 180
conifers, 512
Coniothyrium minitans, 204
conjugation, 799
cost of virulence, 699
cotton, 764
crown defoliation, 502
crucifer black rot, 416
cucumber, 409, 678
Cucumis melo, 339
Cucurbita argyrosperma, 448
cultivar mixture, 723
cultivar ranking, 169
cultural control, 587

D1 (herbicide-binding) protein, 671
DAS-ELISA, 621
Daucus carota, 665
decision support system, 169
defence responses, 227
density dependence, 287
detached leaf dip assay, 512
detection, 416
diagnosis, 1
diagnostic protocols, 161
Diaporthales, 460
disease assessment, 169
disease epidemic, 265
disease forecasting, 347, 357
disease incidence, 108, 535
disease pattern, 347
disease resistance, 483
disease resistance screening, 764
disease severity, 535
DNA fingerprinting, 733
dormancy, 535
dot-blot, 29
dsRNA, 196

ELISA, 431
epidemic development, 144
epidemiology, 287, 299, 522
ergot, 749
Erwinia pyrifoliae, 789
ethylene, 773

© 2005 BSPP

Euphorbia heterophylla, 448
evolutionary diversity, 8
extracellular enzymes, 15

field disease assessment, 364
fire blight, 108
fireblight, 789
fitness cost, 723
fluorescence, 671
foliage, 512
forest species, 512
frosty pod of cocoa, 483
fungi, 189
Fusarium circinatum, 46
Fusarium culmorum, 156
Fusarium graminearum, 156
fusarium head blight, 299
Fusarium langsethiae, 299
Fusarium oxysporum, 53
Fusarium oxysporum f.sp. *asparagi*, 227
Fusarium oxysporum f.sp. *lycopersici*, 773
Fusarium oxysporum f. sp. *melonis*, 339
fusarium wilt, 339

gas exchange analysis, 431
geminivirus, 448
gene dosage, 93
gene sequencing, 789
genetic diversity, 46, 125, 189
genome sequencing, 615
genotype × environment, 364
gentisic acid, 441
Glycine max, 53
grey mould, 317

Himalayas, 36
histology, 535
histopathology, 657
HMA, 625
host growth, 275
host resistance, 287
host specificity, 740
host–parasite relationships, 657
hrp mutants, 781
hydrogen peroxide, 317
Hyparrhenia, 749
hypersensitive reaction, 476
hypersensitive response, 82, 781
hypersensitivity, 308
hypovirulence, 196

IGS, 36
ilarviruses, 29
in vivo pathogenicity, 156
incompatible interaction, 308
incubation, 347
induced resistance, 678
induced systemic resistance, 773
infection, 82, 357
insecticides, 587
integrated disease management, 401
integrated management, 275
interspecific hybrids, 549
invasive plant, 275
isolate-specific resistance, 134

ISSR-PCR, 650
ITS, 36
ITS, 325

late blight, 373, 740
latent infections, 22
latent period, 287
leaf cutter bees, 204
leaf spot, 275
lenticels, 74
lesion development, 144
lesion growth, 265
lime, 393
Loudetia, 749
Lycopersicon esculentum, 448

Malus, 125
Malus domestica, 116
Manihot esculenta, 587
Medicago sativa, 204
Meloidogyne exigua, 471, 476
Meloidogyne paranaensis, 471
Meloidogyne spp., 93
metalaxyl resistance, 733, 740
Mi resistance gene, 93
Miconia rubiginosa, 460
Miconia theaezans, 460
Microdochium nivale, 299
microsatellite markers, 189
microsatellites, 650
migration, 733
mild-strain protection, 587
mitochondrial SSU rDNA, 53
mixed viral infection, 448
MNSV, 615
molecular markers, 46
mollicutes, 625
moniliasis, 483
monoclonal antibodies, 100
mortality, 357
multiplex PCR, 416
Mycosphaerella graminicola, 134
mycotoxins, 299
Myzus persicae, 100

negative binomial distribution, 471
nematode reproduction, 657
nematodes, 471
Nicotiana tabacum, 650
nitrogen uptake, 491
nonpathogenic isolates, 227
nuclear rDNA, 53

oak decline, 502
onion white rot, 212
oomycete, 522
oospores, 373

parasitic angiosperm, 650
partial resistance, 169
pathogen fitness, 265
pathogenesis-related proteins, 678
pathogenicity, 15, 196
pathotypes, 180, 549
PCR, 373, 643, 686, 764

PCR detection, 66
PCR inhibitors, 1
pea, 409
peanut, 401
pear, 108
pear scab, 357
perennation, 535
peroxidase, 317
Phaeoisariopsis personata, 401
Phialophora gregata, 180
photosynthesis, 431
phylogenetic analysis, 621
phylogenetic species, 36
physiological races, 116, 125
Phytophthora infestans, 723
Phytophthora quercina, 502
phytoplasma, 634
phytoplasmas, 8, 625
phytosanitation, 587
phytotoxin, 671
Pierce's disease, 1
Pinus halepensis, 331
Pinus spp., 46
pitch canker, 46
plant growth stage, 22
plant metabolic inhibitors, 409
Podosphaera leucotricha, 116, 125
Podosphaera xanthii, 678
polerovirus, 100
population dynamics, 212
population genetics, 522, 733
population structure, 189
populations, 686
postharvest diseases, 74
potato, 373
potato and tomato late blight, 733
potato late blight, 169, 364
Potato virus Y^{NTN}, 441
potato, 441
powdery mildew, 125, 308
PR proteins, 773
precipitation, 502
principal component analysis, 364
procymidone, 393
progeny testing, 764
Pseudocercosporella herpotrichoides, 144
Pseudomonas aeruginosa, 219
PSII, 671
Puccinia striiformis, 699
Puccinia triticina, 287
Pyrenophora teres f. *maculata*, 66
Pyrenophora teres f. *teres*, 66
Pyrus spp., 357

quantitative PCR analysis, 156
quantitative resistance, 364
quarantine organism, 643
quarantine pathogens, 161
quince, 108

race, 331
race-nonspecific resistance, 364
RAPD, 325
RAPDs, 331
rDNA, 325
reactive oxygen species, 308
real-time PCR, 1, 22
recombination, 621
resistance, 82, 265, 476, 650
resistance breeding, 483
resistance durability, 723
resistant varieties, 587
RFLP, 36
Rhizoctonia carotae, 665
rhizosphere, 799
riboprobe, 29
RNA-RNA blot hybridization, 196
root and stem rot, 339
root length density, 491
root-knot nematode, 476
Rosaceae, 789
rose breeding, 82
rp operon, 8
RT–PCA, 100
RT–PCR, 621
RT–PCR–RFLP, 615
Rubus, 789

Saccharosydne saccharivora, 634
salicylic acid, 227, 441
SCAR markers, 66
scleroderris canker, 331
sclerotial germination, 393
Sclerotinia sclerotiorum, 204, 665
Sclerotium cepivorum, 212
seedborne disease, 416, 643
septoria tritici blotch, 134
sesame, 409
silicon, 678
silthiofam, 686
soil chemistry, 502
soil moisture, 22
soil type, 22
soil water content, 491
soilborne disease, 686
soil-free growing system, 219
Solanum tuberosum, 383, 441, 657
sorghum, 749
soybean, 180
soybean bacterial pustule, 409
soybean root rot, 53
species coexistence, 144
Sphaerotheca fuliginea, 678
SSR, 522
Stb6, 134
storage boxes, 665
storage diseases of carrot, 665
Streptomyces europaeiscabiei, 383
Streptomyces reticuliscabiei, 383
Streptomyces scabies, 383
streptomycin, 108
sudden oak death, 512
sugarcane, 634
superoxide, 317
superoxide dismutase, 317
susceptibility, 476
systemic acquired resistance, 227

take-all, 491, 686
taxonomy, 161

temperature, 22, 502
temperature change, 156
threshold level, 657
Ti plasmid, 799
tiger lily, 621
tissue immunoblot, 615
tissue-printing, 29
tobacco, 409
tobacco cultivars, 650
tolerance, 431
tomato, 29, 409
tomato genotypes, 93
trade-off, 265
transformation, 799
transmission studies, 100
transpiration, 431
Trichoderma atroviride, 204, 212
Triticum aestivum, 287, 491, 699
tuf gene, 8
type III secretion system, 781

UP-PCR band profile, 212

VCG, 339
vector transmission, 634
vegetative compatibility, 549
virulence, 93, 116, 196
virus strains, 615
virus variability, 621
Vitis vinifera, 522, 535

water uptake, 491
water-use efficiency, 431
Web-Blight, 169
wheat, 134, 308, 686
wheat yellow rust, 699
whitefly vector, 587
wild Solanaceae, 740
wilt diseases, 549
wounds, 74

Xanthomonas oryzae pv. *oryzae*, 15
xpsE, 15

yellow leaf syndrome, 634

zoosporic pathogens, 219